arXiv:1008.5207v1 [math.NT] 31 Aug 2010

ON THE FERMIONIC p-ADIC INTEGRAL REPRESENTATION OF BERNSTEIN POLYNOMIALS ASSOCIATED WITH EULER NUMBERS AND POLYNOMIALS

T. KIM, J. CHOI, Y.-H. KIM, AND C. S. RYOO

Abstract The purpose of this paper is to give some properties of several Bernstein type polynomials to represent the fermionic p-adic integral on $\mathbb{Z}_p$. From these properties, we derive some interesting identities on the Euler numbers and polynomials.

1. Introduction

Throughout this paper, let p be an add prime number. The symbol $\mathbb{Z}_p$, $\mathbb{Q}_p$ and $\mathbb{C}_p$ denote the ring of p-adic integers, the field of p-adic numbers and the field of p-adic completion of the algebraic closure of $\mathbb{Q}_p$, respectively (see [1-14]).

Let $\mathbb{N}$ be the set of natural numbers and $\mathbb{Z}_+ = \mathbb{N} \cup \{0\}$. Let ν_p be the normalized exponential valuation of $\mathbb{C}_p$ with $|p|_p = p^{-\nu_p(p)} = \frac{1}{p}$. Note that $\mathbb{Z}_p = \{x \mid |x|_p \leq 1\} = \varprojlim_N \mathbb{Z}/p^N\mathbb{Z}_p$.

When one talks of q-extension, q is variously consiered as an indeterminate, a complex number $q \in \mathbb{C}$ or p-adic number $q \in \mathbb{C}_p$. If $q \in \mathbb{C}$, we normally assume $|q| < 1$ and if $q \in \mathbb{C}_p$, we always assume $|1 - q|_p < 1$.

We say that f is uniformly differentiable function at a point $a \in \mathbb{Z}$ and write $f \in UD(\mathbb{Z}_p)$, if the difference quotient $F_f(x, y) = \frac{f(x)-f(y)}{x-y}$ have a limit $f'(a)$ as $(x, y) \to (a, a)$. For $f \in UD(\mathbb{Z}_p)$, the fermionic p-adic q-integral on $\mathbb{Z}_p$ is defined as

$$I_{-q}(f) = \int_{\mathbb{Z}_p} f(x) d\mu_{-q}(x) = \lim_{N\to\infty} \frac{1+q}{1+q^{p^N}} \sum_{x=0}^{p^N-1} f(x)(-q)^x, \quad \text{(see [7]).} \tag{1}$$

In the special case $q = 1$ in (1), the integral

$$I_{-1}(f) = \int_{\mathbb{Z}_p} f(x) d\mu_{-1}(x), \tag{2}$$

is called the fermionic p-adic invariant integral on $\mathbb{Z}_p$ (see [12]). From (2), we note

$$I_{-1}(f_1) = -I_{-1}(f) + 2f(0), \tag{3}$$

where $f_1(x) = f(x + 1)$.

2000 Mathematics Subject Classification : 11B68, 11B73, 41A30.

Key words and phrases : Bernstein polynomial, Euler numbers and polynomials, fermionic p-adic intergal.

Moreover, for $n \in \mathbb{N}$, let $f_n(x) = f(x+n)$. Then we note that

$$I_{-1}(f_n) = (-1)^n I_{-1}(f) + 2\sum_{l=0}^{n-1}(-1)^{n-1-l}f(l). \tag{4}$$

It is well known that the Euler polynomials are defined by

$$\frac{2}{e^t+1}e^{xt} = \sum_{n=0}^{\infty} E_n(x)\frac{t}{n!}, \quad \text{(see [1-14])}. \tag{5}$$

In the special case, $x = 0$, $E_n(0) = E_n$ are called the n-th Euler numbers.

Let $f(x) = e^{tx}$. Then, by (3), (4) and (5), we see that

$$\int_{\mathbb{Z}_p} e^{(x+y)t}d\mu_{-1}(y) = \frac{2}{e^t+1}e^{xt} = \sum_{n=0}^{\infty} E_n(x)\frac{t^n}{n!}. \tag{6}$$

Let $C[0,1]$ denote the set of continuous functions on $[0,1]$. For $f \in C[0,1]$, Bernstein introduced the following well-known linear positive operator in the field of real numbers $\mathbb{R}$:

$$\mathbb{B}_n(f:x) = \sum_{k=0}^{n} f(\frac{k}{n})\binom{n}{k}x^k(1-x)^{n-k} = \sum_{k=0}^{n} f(\frac{k}{n})B_{k,n}(x), \tag{7}$$

where $\binom{n}{k} = \frac{n(n-1)\cdots(n-k+1)}{k!} = \frac{n!}{k!(n-k)!}$ (see [1, 2, 5, 9, 10, 14]). Here, $\mathbb{B}_n(f:x)$ is called the Bernstein operator of order n for f.

For $k, n \in \mathbb{Z}_+$, the Bernstein polynomial of degree n is defined by

$$B_{k,n} = \binom{n}{k}x^k(1-x)^{n-k}, \quad \text{for } x \in [0,1]. \tag{8}$$

For example, $B_{0,1}(x) = 1-x$, $B_{1,1}(x) = x$, $B_{0,2}(x) = (1-x)^2$, $B_{1,2}(x) = 2x-2x^2$, $B_{2,2}(x) = x^2, \ldots$, and $B_{k,n}(x) = 0$ for $n < k$, $B_{k,n}(x) = B_{n-k,n}(1-x)$.

In this paper, we study the properties of Bernstein polynomials in the p-adic number field. For $f \in UD(\mathbb{Z}_p)$, we give some properties of several type Bernstein polynomials to represent the fermionic p-adic invariant integral on $\mathbb{Z}_p$. From those properties, we derive some interesting identities on the Euler polynomials.

2. Fermionic p-adic integral representation of Bernstein polynomials

By (5) and (6), we see that

$$\frac{2}{e^t+1}e^{(1-x)t} = \sum_{n=0}^{\infty} E_n(1-x)\frac{t^n}{n!}. \tag{9}$$

We also have that

$$\frac{2}{e^t+1}e^{(1-x)t} = \frac{2}{1+e^{-t}}e^{-xt} = \sum_{n=0}^{\infty} E_n(x)\frac{(-1)^n}{n!}t^n. \tag{10}$$

From (9) and (10), we note tht $E_n(1-x) = (-1)^n E_n(x)$. It is easy to show that

$$E_n(2) = 2 - \sum_{l=0}^{n}\binom{n}{l}E_l = 2 + E_n, \quad \text{for } n > 0. \tag{11}$$

By (5), (6), (9), (10) and (11), we see that for $n > 0$,

$$\begin{aligned}\int_{\mathbb{Z}_p}(1-x)^n d\mu_{-1}(x) &= (-1)^n\int_{\mathbb{Z}_p}(x-1)^n d\mu_{-1}(x)\\ &= \int_{\mathbb{Z}_p}(x+2)^n d\mu_{-1}(x)\\ &= 2+\int_{\mathbb{Z}_p}x^n d\mu_{-1}(x).\end{aligned}$$

Therefore, we obtain the following theorem.

Theorem 1. *For $n \in \mathbb{N}$, we have*

$$\int_{\mathbb{Z}_p}(1-x)^n d\mu_{-1}(x) = 2+\int_{\mathbb{Z}_p}x^n d\mu_{-1}(x).$$

Theorem 1 is important to derive our main result in this paper.

Taking the fermionic p-adic integral on $\mathbb{Z}_p$ for one Bernstein polynomial in (8), we get

$$\begin{aligned}\int_{\mathbb{Z}_p}B_{k,n}(x)d\mu_{-1}(x) &= \int_{\mathbb{Z}_p}\binom{n}{k}x^k(1-x)^{n-k}d\mu_{-1}(x)\\ &= \binom{n}{k}\sum_{j=0}^{n-k}\binom{n-k}{j}(-1)^{n-k-j}\int_{\mathbb{Z}_p}x^{n-j}d\mu_{-1}(x)\\ &= \binom{n}{k}\sum_{j=0}^{n-k}\binom{n-k}{j}(-1)^{n-k-j}E_{n-j}\\ &= \binom{n}{k}\sum_{j=0}^{n-k}\binom{n-k}{j}(-1)^{j}E_{k+j}.\end{aligned}$$

Therefore, we obtain the following proposition.

Proposition 2. *For $k, n \in \mathbb{Z}_+$, we have*

$$\int_{\mathbb{Z}_p}B_{k,n}(x)d\mu_{-1}(x) = \binom{n}{k}\sum_{j=0}^{n-k}\binom{n-k}{j}(-1)^{j}E_{k+j}.$$

It is known that $B_{k,n}(x) = B_{n-k,n}(1-x)$. Thus, we have

$$\begin{aligned}\int_{\mathbb{Z}_p}B_{k,n}(x)d\mu_{-1}(x) &= \int_{\mathbb{Z}_p}B_{n-k,n}(1-x)d\mu_{-1}(x) \qquad (12)\\ &= \binom{n}{n-k}\sum_{j=0}^{k}\binom{k}{j}(-1)^{k-j}\int_{\mathbb{Z}_p}(1-x)^{n-j}d\mu_{-1}(x).\end{aligned}$$

By (12) and Theorem 1, we see that for $n > k$,

$$\begin{aligned}\int_{\mathbb{Z}_p} B_{k,n}(x)d\mu_{-1}(x) &= \binom{n}{k}\sum_{j=0}^{k}\binom{k}{j}(-1)^{k-j}(2+\int_{\mathbb{Z}_p} x^{n-j}d\mu_{-1}(x)) \\ &= \binom{n}{k}\sum_{j=0}^{k}\binom{k}{j}(-1)^{k-j}(2+E_{n-j}) \qquad (13) \\ &= \begin{cases} 2+E_n & \text{if } k=0, \\ \binom{n}{k}\sum_{j=0}^{k}\binom{k}{j}(-1)^{k-j}E_{n-j} & \text{if } k>0. \end{cases}\end{aligned}$$

From (13), we obtain the following theorem.

Theorem 3. *For $n, k \in \mathbb{Z}_+$ with $n > k$, we have*

$$\int_{\mathbb{Z}_p} B_{k,n}(x)d\mu_{-1}(x) = \begin{cases} 2+E_n & \textit{if } k=0, \\ \binom{n}{k}\sum_{j=0}^{k}\binom{k}{j}(-1)^{k-j}E_{n-j} & \textit{if } k>0. \end{cases}$$

By Proposition 2 and Theorem 3, we obtain the following corollary.

Corollary 4. *For $n, k \in \mathbb{Z}_+$ with $n > k$, we have*

$$\sum_{j=0}^{n-k}\binom{n-k}{j}(-1)^j E_{k+j} = \begin{cases} 2+E_n & \textit{if } k=0, \\ \sum_{j=0}^{k}\binom{k}{j}(-1)^{k-j}E_{n-j} & \textit{if } k>0. \end{cases}$$

For $m, n, k \in \mathbb{Z}_+$ with $m + n > 2k$, fermionic p-adic invariant integral for multiplication of two Berstein polynomials on $\mathbb{Z}_p$ can be given by the following relation.

$$\begin{aligned}&\int_{\mathbb{Z}_p} B_{k,n}(x)B_{k,m}(x)d\mu_{-1}(x) \\ &= \int_{\mathbb{Z}_p}\binom{n}{k}x^k(1-x)^{n-k}\binom{m}{k}x^k(1-x)^{m-k}d\mu_{-1}(x) \\ &= \binom{n}{k}\binom{m}{k}\int_{\mathbb{Z}_p} x^{2k}(1-x)^{n+m-2k}d\mu_{-1}(x) \\ &= \binom{n}{k}\binom{m}{k}\sum_{j=0}^{2k}\binom{2k}{j}(-1)^{j+2k}\int_{\mathbb{Z}_p}(1-x)^{n+m-j}d\mu_{-1}(x) \\ &= \binom{n}{k}\binom{m}{k}\sum_{j=0}^{2k}\binom{2k}{j}(-1)^{j+2k}(2+\int_{\mathbb{Z}_p} x^{n+m-j}d\mu_{-1}(x)) \\ &= \begin{cases} 2+E_{n+m} & \text{if } k=0, \\ \binom{n}{k}\binom{m}{k}\sum_{j=0}^{2k}\binom{2k}{j}(-1)^{j+2k}E_{n+m-j} & \text{if } k>0. \end{cases}\end{aligned}$$

Therefore, we obtain the following theorem.

Theorem 5. *For $m, n, k \in \mathbb{Z}_+$ with $m+n>2k$, we have*

$$\begin{aligned}&\int_{\mathbb{Z}_p} B_{k,n}(x)B_{k,m}(x)d\mu_{-1}(x)\\ &\quad= \begin{cases} 2+E_{n+m} & \text{if } k=0,\\ \binom{n}{k}\binom{m}{k}\sum_{j=0}^{2k}\binom{2k}{j}(-1)^{j+2k}E_{n+m-j} & \text{if } k>0.\end{cases}\end{aligned}$$

For $m, n, k \in \mathbb{Z}_+$, we have

$$\begin{aligned}&\int_{\mathbb{Z}_p} B_{k,n}(x)B_{k,m}(x)d\mu_{-1}(x)\\ &\quad=\binom{n}{k}\binom{m}{k}\int_{\mathbb{Z}_p} x^{2k}(1-x)^{n+m-2k}d\mu_{-1}(x)\\ &\quad=\binom{n}{k}\binom{m}{k}\sum_{j=0}^{n+m-2k}\binom{n+m-2k}{j}(-1)^j\int_{\mathbb{Z}_p} x^{j+2k}d\mu_{-1}(x)\\ &\quad=\binom{n}{k}\binom{m}{k}\sum_{j=0}^{n+m-2k}\binom{n+m-2k}{j}(-1)^j E_{j+2k}.\end{aligned}$$

Thus, we obtain the following proposition.

Proposition 6. *For $m, n, k \in \mathbb{Z}_+$, we have*

$$\int_{\mathbb{Z}_p} B_{k,n}(x)B_{k,m}(x)d\mu_{-1}(x)=\binom{n}{k}\binom{m}{k}\sum_{j=0}^{n+m-2k}\binom{n+m-2k}{j}(-1)^j E_{j+2k}.$$

By Theorem 5 and Proposition 6, we obtain the following corollary.

Corollary 7. *For $m, n, k \in \mathbb{Z}_+$ with $m+n>2k$, we have*

$$\begin{aligned}&\sum_{j=0}^{n+m-2k}\binom{n+m-2k}{j}(-1)^j E_{j+2k}\\ &\quad= \begin{cases} 2+E_{n+m} & \text{if } k=0,\\ \sum_{j=0}^{2k}\binom{2k}{j}(-1)^{j+2k}E_{n+m-j} & \text{if } k>0.\end{cases}\end{aligned}$$

In the same manner, multiplication of three Bernstein polynomials can be given by the following relation :

$$\begin{aligned}&\int_{\mathbb{Z}_p} B_{k,n}(x)B_{k,m}(x)B_{k,s}(x)d\mu_{-1}(x)\\ &=\binom{n}{k}\binom{m}{k}\binom{s}{k}\sum_{j=0}^{n+m+s-3k}\binom{n+m+s-3k}{j}(-1)^j\int_{\mathbb{Z}_p} x^{j+3k}d\mu_{-1}(x) \qquad (14)\\ &=\binom{n}{k}\binom{m}{k}\binom{s}{k}\sum_{j=0}^{n+m+s-3k}\binom{n+m+s-3k}{j}(-1)^j E_{j+3k},\end{aligned}$$

where $m, n, s, k \in \mathbb{Z}_+$ with $m+n+s>3k$.

For $m, n, s, k \in \mathbb{Z}_+$ with $m+n+s > 3k$, by the symmetry of Bernstein polynomials, we see that

$$\begin{aligned}
&\int_{\mathbb{Z}_p} B_{k,n}(x)B_{k,m}(x)B_{k,s}(x)d\mu_{-1}(x)\\
&\quad= \binom{n}{k}\binom{m}{k}\binom{s}{k}\sum_{j=0}^{3k}\binom{3k}{j}(-1)^{3k-j}\int_{\mathbb{Z}_p}(1-x)^{n+m+s-j}d\mu_{-1}(x)\\
&\quad= \binom{n}{k}\binom{m}{k}\binom{s}{k}\sum_{j=0}^{3k}\binom{3k}{j}(-1)^{3k-j}(2+\int_{\mathbb{Z}_p}x^{n+m+s-j}\mu_{-1}(x))\\
&\quad= \begin{cases} 2+E_{n+m+s} & \text{if } k=0,\\ \binom{n}{k}\binom{m}{k}\binom{s}{k}\sum_{j=0}^{3k}\binom{3k}{j}(-1)^{3k-j}E_{n+m+s-j} & \text{if } k>0. \end{cases}
\end{aligned}$$

Therefore, we obtain the following theorem.

Theorem 8. *For $m, n, s, k \in \mathbb{Z}_+$ with $m+n+s > 3k$, we have*

$$\begin{aligned}
&\int_{\mathbb{Z}_p} B_{k,n}(x)B_{k,m}(x)B_{k,s}(x)d\mu_{-1}(x)\\
&\quad= \begin{cases} 2+E_{n+m+s} & \text{if } k=0,\\ \binom{n}{k}\binom{m}{k}\binom{s}{k}\sum_{j=0}^{3k}\binom{3k}{j}(-1)^{3k-j}E_{n+m+s-j} & \text{if } k>0. \end{cases}
\end{aligned}$$

By (14) and Theorem 8, we obtain the following corollary.

Corollary 9. *For $m, n, s, k \in \mathbb{Z}_+$ with $m+n+s > 3k$, we have*

$$\begin{aligned}
&\sum_{j=0}^{n+m+s-3k}\binom{n+m+s-3k}{j}(-1)^j E_{j+3k}\\
&\quad= \begin{cases} 2+E_{n+m+s} & \text{if } k=0,\\ \sum_{j=0}^{3k}\binom{3k}{j}(-1)^{3k-j}E_{n+m+s-j} & \text{if } k>0. \end{cases}
\end{aligned}$$

Using the above theorems and mathematical induction, we obtain the following theorem.

Theorem 10. *Let $s \in \mathbb{N}$. For $n_1, n_2, \ldots, n_s, k \in \mathbb{Z}_+$ with $n_1+n_2+\cdots+n_s > sk$, the multiplication of the sequence of Bernstein polynomials $B_{k,n_1}(x), \ldots, B_{k,n_s}(x)$ with different degrees under fermionic p-adic invariant integral on $\mathbb{Z}_p$ can be given as*

$$\begin{aligned}
&\int_{\mathbb{Z}_p}\left(\prod_{i=1}^{s}B_{k,n_i}(x)\right)d\mu_{-1}(x)\\
&\quad= \begin{cases} 2+E_{n_1+n_2+\cdots+n_s} & \text{if } k=0,\\ \left(\prod_{i=1}^{s}\binom{n_i}{k}\right)\sum_{j=0}^{sk}\binom{sk}{j}(-1)^{sk-j}E_{n_1+n_2+\cdots+n_s-j} & \text{if } k>0. \end{cases}
\end{aligned}$$

We also easily see that

$$\int_{\mathbb{Z}_p}\left(\prod_{i=1}^{s}B_{k,n_i}(x)\right)d\mu_{-1}(x) \tag{15}$$
$$=\left(\prod_{i=1}^{s}\binom{n_i}{k}\right)\sum_{j=0}^{n_1+\cdots+n_s-sk}\binom{n_1+\cdots+n_s-sk}{j}(-1)^jE_{j+sk}.$$

By Theorem 10 and (15), we obtain the following corollary.

Corollary 11. *Let $s\in\mathbb{N}$. For $n_1,n_2,\ldots,n_s,k\in\mathbb{Z}_+$ with $n_1+n_2+\cdots+n_s>sk$, we have*

$$\sum_{j=0}^{n_1+\cdots+n_s-sk}\binom{n_1+\cdots+n_s-sk}{j}(-1)^jE_{j+sk}$$
$$=\begin{cases}2+E_{n_1+n_2+\cdots+n_s} & \textit{if}\ \ k=0,\\ \sum_{j=0}^{sk}\binom{sk}{j}(-1)^{sk-j}E_{n_1+n_2+\cdots+n_s-j} & \textit{if}\ \ k>0.\end{cases}$$

Let $m_1,\ldots,m_s,n_1,\ldots,n_s,k\in\mathbb{Z}_+$ with $m_1n_1+\cdots+m_sn_s>(m_1+\cdots+m_s)k$. By the definition of $B_{k,n_s}^{m_s}(x)$, we easily get

$$\int_{\mathbb{Z}_p}\left(\prod_{i=1}^{s}B_{k,n_i}^{m_i}(x)\right)d\mu_{-1}(x)$$
$$=\left(\prod_{i=1}^{s}\binom{n_i}{k}^{m_i}\right)\sum_{j=0}^{k\sum_{i=1}^{s}m_i}(-1)^{k\sum_{i=1}^{s}m_i-j}\int_{\mathbb{Z}_p}(1-x)^{\sum_{i=1}^{s}n_im_i-j}d\mu_{-1}(x)$$
$$=\left(\prod_{i=1}^{s}\binom{n_i}{k}^{m_i}\right)\sum_{j=0}^{k\sum_{i=1}^{s}m_i}\binom{k\sum_{i=1}^{s}m_i}{j}(-1)^{k\sum_{i=1}^{s}m_i-j}(2+E_{\sum_{i=1}^{s}n_im_i-j})$$
$$=\begin{cases}2+E_{m_1n_1+\cdots+m_sn_s} & \text{if}\ \ k=0,\\ \left(\prod_{i=1}^{s}\binom{n_i}{k}^{m_i}\right)\sum_{j=0}^{k\sum_{i=1}^{s}m_i}\binom{k\sum_{i=1}^{s}m_i}{j}(-1)^{k\sum_{i=1}^{s}m_i-j}E_{\sum_{i=1}^{s}n_im_i-j} & \text{if}\ \ k>0.\end{cases}$$

Therefore, we obtain the following theorem.

Theorem 12. *Let $s\in\mathbb{N}$. For $m_1,\ldots,m_s,n_1,\ldots,n_s,k\in\mathbb{Z}_+$ with $m_1n_1+\cdots+m_sn_s>(m_1+\cdots+m_s)k$, we have*

$$\int_{\mathbb{Z}_p}\left(\prod_{i=1}^{s}B_{k,n_i}^{m_i}(x)\right)d\mu_{-1}(x)$$
$$=\begin{cases}2+E_{m_1n_1+\cdots+m_sn_s} & \textit{if}\ \ k=0,\\ \left(\prod_{i=1}^{s}\binom{n_i}{k}^{m_i}\right)\sum_{j=0}^{k\sum_{i=1}^{s}m_i}\binom{k\sum_{i=1}^{s}m_i}{j}(-1)^{k\sum_{i=1}^{s}m_i-j}E_{\sum_{i=1}^{s}n_im_i-j} & \textit{if}\ \ k>0.\end{cases}$$

By simple calculation, we easily get

$$\begin{aligned}
&\int_{\mathbb{Z}_p}\left(\prod_{i=1}^{s} B_{k,n_i}^{m_i}(x)\right) d\mu_{-1}(x)\\
&=\left(\prod_{i=1}^{s}\binom{n_i}{k}^{m_i}\right)\sum_{j=0}^{\sum_{i=1}^{s} n_i m_i - k\sum_{i=1}^{s} m_i}\binom{\sum_{i=1}^{s} n_i m_i - k\sum_{i=1}^{s} m_i}{j}(-1)^j E_{k\sum_{i=1}^{s} m_i - j},
\end{aligned}$$

where $m_1, \ldots, m_s, n_1, \ldots, n_s, k \in \mathbb{Z}_+$ for $s \in \mathbb{N}$. By Theorem 12 and (16), we obtain the following corollary.

Corollary 13. *Let $s \in \mathbb{N}$. For $m_1, \ldots, m_s, n_1, \ldots, n_s, k \in \mathbb{Z}_+$ with $m_1 n_1 + \cdots + m_s n_s > (m_1 + \cdots + m_s)k$, we have*

$$\begin{aligned}
&\sum_{j=0}^{\sum_{i=1}^{s} n_i m_i - k\sum_{i=1}^{s} m_i}\binom{\sum_{i=1}^{s} n_i m_i - k\sum_{i=1}^{s} m_i}{j}(-1)^j E_{k\sum_{i=1}^{s} m_i - j}\\
&=\begin{cases} 2 + E_{m_1 n_1 + \cdots + m_s n_s} & \text{if } \ k = 0,\\ \sum_{j=0}^{k\sum_{i=1}^{s} m_i}\binom{k\sum_{i=1}^{s} m_i}{j}(-1)^{k\sum_{i=1}^{s} m_i - j} E_{\sum_{i=1}^{s} n_i m_i - j} & \text{if } \ k > 0.\end{cases}
\end{aligned}$$

The fermionic p-adic invariant integral of multiplication of $(n+1)$ Bernstein polynomials, the n-th degree Bernstein polynomials $B_{i,n}(x)$ with $i = 0, 1, \ldots, n$ and with multiplicity $m_0, m_1, \ldots, m_n$ on $\mathbb{Z}_p$, respectively, can be given by

$$\begin{aligned}
&\int_{\mathbb{Z}_p}\left(\prod_{i=0}^{n} B_{i,n}^{m_i}(x)\right) d\mu_{-1}(x)\\
&=\left(\prod_{i=0}^{n}\binom{n}{i}^{m_i}\right)\int_{\mathbb{Z}_p} x^{\sum_{i=1}^{n} i m_i}(1-x)^{n\sum_{i=0}^{n} m_i - \sum_{i=1}^{n} i m_i} d\mu_{-1}(x)\\
&=\frac{\left(\prod_{i=1}^{n}\binom{n}{i}^{m_i}\right)}{\binom{n\sum_{i=0}^{n} m_i}{\sum_{i=1}^{n} i m_i}}\int_{\mathbb{Z}_p} B_{\sum_{i=1}^{n} i m_i,\, n\sum_{i=0}^{n} m_i}(x) d\mu_{-1}(x),
\end{aligned}$$

where $m_0, m_1, \ldots, m_n \in \mathbb{Z}_+$ with $n \in \mathbb{Z}_+$.

Assume that $n m_0 + n m_1 + \cdots + n m_n > m_1 + 2m_2 + \cdots + n m_n$. Then we have

$$\begin{aligned}
&\int_{\mathbb{Z}_p}\left(\prod_{i=0}^{n} B_{i,n}^{m_i}(x)\right) d\mu_{-1}(x)\\
&=\begin{cases} 2 + E_{n m_0 + n m_1 + \cdots + n m_n} & \text{if } \ \sum_{i=1}^{n} i m_i = 0,\\ \left(\prod_{i=0}^{n}\binom{n}{i}^{m_i}\right)\sum_{j=0}^{\sum_{i=1}^{n} i m_i}\binom{\sum_{i=1}^{n} i m_i}{j}(-1)^{\sum_{i=1}^{n} i m_i - j} E_{n\sum_{i=0}^{n} m_i - \sum_{i=1}^{n} i m_i} & \\ & \text{if } \ \sum_{i=1}^{n} i m_i > 0.\end{cases}
\end{aligned}$$

Therefore, we obtain the following theorem.

Theorem 14. *Let $n \in \mathbb{Z}_+$.*
(I) For $m_0, m_1, \ldots, m_n \in \mathbb{Z}_+$ with $n\sum_{i=0}^n m_i > \sum_{i=1}^n im_i$, we have

$$\begin{aligned}
&\int_{\mathbb{Z}_p}\left(\prod_{i=0}^n B_{i,n}^{m_i}(x)\right)d\mu_{-1}(x)\\
&=\begin{cases} 2+E_{nm_0+nm_1+\cdots+nm_n} & \text{if } \sum_{i=1}^n im_i = 0,\\ \left(\prod_{i=0}^n \binom{n}{i}^{m_i}\right)\sum_{j=0}^{\sum_{i=1}^n im_i}\binom{\sum_{i=1}^n im_i}{j}(-1)^{\sum_{i=1}^n im_i-j}E_{n\sum_{i=0}^n m_i-\sum_{i=1}^n im_i} & \\ & \text{if } \sum_{i=1}^n im_i > 0.\end{cases}
\end{aligned}$$

(II) For $m_0, m_1, \ldots, m_n \in \mathbb{Z}_+$, we have

$$\begin{aligned}
&\int_{\mathbb{Z}_p}\left(\prod_{i=0}^n B_{i,n}^{m_i}(x)\right)d\mu_{-1}(x)\\
&=\left(\prod_{i=0}^n \binom{n}{i}^{m_i}\right)\sum_{j=0}^{n\sum_{i=0}^n m_i-\sum_{i=1}^n im_i}\binom{n\sum_{i=0}^n m_i-\sum_{i=1}^n im_i}{j}(-1)^j E_{\sum_{i=1}^n im_i+j}.
\end{aligned}$$

By Theorem 14, we obtain the following corollary.

Corollary 15. *For $n, m_0, m_1, \ldots, m_n \in \mathbb{Z}_+$ with $n\sum_{i=0}^n m_i > \sum_{i=1}^n im_i$, we have*

$$\begin{aligned}
&\sum_{j=0}^{n\sum_{i=0}^n m_i-\sum_{i=1}^n im_i}\binom{n\sum_{i=0}^n m_i-\sum_{i=1}^n im_i}{j}(-1)^j E_{\sum_{i=1}^n im_i+j}\\
&=\begin{cases} 2+E_{nm_0+nm_1+\cdots+nm_n} & \text{if } \sum_{i=1}^n im_i = 0,\\ \sum_{j=0}^{\sum_{i=1}^n im_i}\binom{\sum_{i=1}^n im_i}{j}(-1)^{\sum_{i=1}^n im_i-j}E_{n\sum_{i=0}^n m_i-\sum_{i=1}^n im_i} & \\ & \text{if } \sum_{i=1}^n im_i > 0.\end{cases}
\end{aligned}$$

REFERENCES

[1] M. Acikgoz, S. Araci, *A study on the integral of the product of several type Berstein polynomials*, IST Transaction of Applied Mathematics-Modelling and Simulation, 2010.

[2] S. Bernstein, *Démonstration du théorème de Weierstrass, fondée sur le calcul des probabilities*, Commun. Soc. Math. Kharkow **13** (1912), 1–2.

[3] I. N. Cangul, V. Kurt, H. Ozden, Y. Simsek, *On the higher-order w-q-Genocchi numbers*, Adv. Stud. Contemp. Math. **19** (2009), 39–57.

[4] N. K. Govil, V. Gupta, *Convergence of q-Meyer-König-Zeller-Durrmeyer operators*, Adv. Stud. Contemp. Math. **19** (2009), 97–108.

[5] V. Gupta, T. Kim, J. Choi, Y.-H. Kim, *Generating function for q-Bernstein, q-Meyer-König-Zeller and q-Beta basis*, Automation Computers Applied Mathematics **19** (2001), 7–11.

[6] T. Kim, *q-extension of the Euler formulae and trigonometric functions*, Russ. J. Math. Phys. **14** (2007), 275–278.

[7] T. Kim, *q-Volkenborn integration*, Russ. J. Math. Phys. **9** (2002), 288–299.

[8] T. Kim, *q-Bernoulli numbers and polynomials associated with Gaussian binomial coefficients*, Russ. J. Math. Phys. **15** (2008), 51–57.

[9] T. Kim, J. Choi, Y.-H. Kim, *Some identities on the q-Berstein polynomials, q-Stirling numbers and q-Bernoulli numbers*, Adv. Stud. Contemp. Math. **20** (2010), 335-341

[10] T. Kim, L.-C. Jang, H. Yi, *Note on the modified q-Berstein polynomials*, Discrete Dynamics in Nature and Society (in press), (2010), arXiv 1005.4293.

[11] T. Kim, *Note on the Euler q-zeta functions*, J. Number Theory **129** (2009), 1798–1804.

[12] T. Kim, *Barnes-type multiple q-zeta functions and q-Euler polynomials*, J. Physics A : Math. Theor., **43** (2010), 255201, 11pp.

[13] V. Kurt,*A further symmetric relation on the analogue of the Apostol-Bernoulli and the analogue of the Apostol-Genocchi polynomials*, Appl. Math. Sci. **3** (2009), 53–56.

[14] Y. Simsek, M. Acikgoz, *A new generating function of q-Berstein-type polynomials and their interpolation function*, Abstract and Applied Analysis **2010** (2010), Article ID 769095, 12 pages.

Taekyun Kim. Division of General Education-Mathematics, Kwangwoon University, Seoul 139-701, Republic of Korea,
E-mail address: `tkkim@kw.ac.kr`

Jongsung Choi. Division of General Education-Mathematics, Kwangwoon University, Seoul 139-701, Republic of Korea,
E-mail address: `jeschoi@kw.ac.kr`

Young-Hee Kim. Division of General Education-Mathematics, Kwangwoon University, Seoul 139-701, Republic of Korea,
E-mail address: `yhkim@kw.ac.kr`

Cheon Seoung Ryoo. Department of Mathematics, Hannam University, Daejeon 306-791, Republic of Korea,
E-mail address: `ryoocs@hnu.kr`